

Double Negation Translations as Morphisms

Olivier Hermant

CRI, MINES ParisTech

December 1, 2014

UFRN, Natal

Double-Negation Translations

Double-Negation translations:

- ▶ a shallow way to encode classical logic into intuitionistic
- ▶ Zenon's backend for Dedukti
- ▶ existing translations: Kolmogorov's (1925), Gentzen-Gödel's (1933), Kuroda's (1951), Krivine's (1990), ...

Minimizing the translations:

- ▶ turns more formulæ into themselves;
- ▶ shifts a classical proof into an intuitionistic proof of the *same* formula.

Morphisms

- ▶ A morphism preserves the operations between two structures:

$$\text{Group morphism: } \begin{cases} (\mathbb{Z}, +, 0) & \mapsto & (\mathbb{R}^*, *, 1) \\ h(0) & \rightarrow & 1 \\ h(a + b) & \rightarrow & h(a) * h(b) \end{cases}$$

- ▶ a translation that is a morphism:

$$\begin{aligned} h(P) &= P \\ h(A \wedge B) &= h(A) \wedge h(B) \\ h(A \vee B) &= h(A) \vee h(B) \\ h(A \Rightarrow B) &= h(A) \Rightarrow h(B) \\ h(\forall x A) &= \forall x h(A) \\ h(\exists x A) &= \exists x h(A) \end{aligned}$$

(of course this is the **identity**)

Morphisms

- ▶ A morphism preserves the operations between two structures:

$$\text{Group morphism: } \begin{cases} (\mathbb{Z}, +, 0) \mapsto (\mathbb{R}^*, *, 1) \\ h(0) \rightarrow 1 \\ h(a + b) \rightarrow h(a) * h(b) \end{cases}$$

- ▶ a **more interesting** translation that is a morphism:

$$\begin{aligned} h(P) &= P \\ h(A \wedge B) &= h(A) \wedge_c h(B) \\ h(A \vee B) &= h(A) \vee_c h(B) \\ h(A \Rightarrow B) &= h(A) \Rightarrow_c h(B) \\ h(\forall x A) &= \forall_c x h(A) \\ h(\exists x A) &= \exists_c x h(A) \end{aligned}$$

two kinds of connectives: the **classical** and the **intuitionistic** ones.

Morphisms

- ▶ A morphism preserves the operations between two structures:

$$\text{Group morphism: } \begin{cases} (\mathbb{Z}, +, 0) & \mapsto & (\mathbb{R}^*, *, 1) \\ h(0) & \rightarrow & 1 \\ h(a + b) & \rightarrow & h(a) * h(b) \end{cases}$$

- ▶ a **more interesting** translation that is a morphism:

$$\begin{aligned} h(P) &= P \\ h(A \wedge B) &= h(A) \wedge_c h(B) \\ h(A \vee B) &= h(A) \vee_c h(B) \\ h(A \Rightarrow B) &= h(A) \Rightarrow_c h(B) \\ h(\forall x A) &= \forall_c x h(A) \\ h(\exists x A) &= \exists_c x h(A) \end{aligned}$$

two kinds of connectives: the **classical** and the **intuitionistic** ones.

- ▶ Design a **unified** logic, where we can reason both classically and intuitionistically:

$$\frac{\Gamma \vdash A}{\Gamma \vdash A \vee_i B}$$

$$\frac{\text{strange premises}}{\Gamma \vdash A \vee_c B}$$

Translations that are Morphisms

- ▶ None of the previous translations is a morphism.
- ▶ Dowek has shown one, it is very verbose.
- ▶ We make it lighter.

Plan:

- 1 Classical and Intuitionistic Logic
- 2 Sequent Calculus
- 3 Double Negation Translations
- 4 Morphisms

Classical vs. Intuitionistic

- ▶ The principle of excluded-middle. Should

$$A \vee \neg A$$

be provable ? Yes or no ?

- ▶ **Yes**. This is what is called **classical** logic.
- ▶ Wait a minute !

The Drinker's Principle

In a bar, there is somebody such that, if he drinks, then everybody drinks.

Two Irrationals

There exists $i_1, i_2 \in \mathbb{R} \setminus \mathbb{Q}$ such that $i_1^{i_2} \in \mathbb{Q}$.

A Manicchean World

You are with us, or against us.

Rashomon (A. Kurosawa).

Classical vs. Intuitionistic

- ▶ The principle of excluded-middle. Should

$$A \vee \neg A$$

be provable ? Yes or no ?

- ▶ **No**. This is the **constructivist** school (Brouwer, Heyting, Kolmogorov).
- ▶ Intuitionistic logic is one of those branches. It features the BHK interpretation of proofs:

Witness Property

A proof of $\exists xA$ (in the empty context) gives a **witness** t for the property A .

Disjunction Property

A proof of $A \vee B$ (in the empty context) reduces eventually **either** to a proof of A , **or** to a proof of B .

The Classical Sequent Calculus (LK)

$$\frac{}{\Gamma, A \vdash A, \Delta} \text{ax}$$

$$\frac{\Gamma, A, B \vdash \Delta}{\Gamma, A \wedge B \vdash \Delta} \wedge_L$$

$$\frac{\Gamma \vdash A, \Delta \quad \Gamma \vdash B, \Delta}{\Gamma \vdash A \wedge B, \Delta} \wedge_R$$

$$\frac{\Gamma, A \vdash \Delta \quad \Gamma, B \vdash \Delta}{\Gamma, A \vee B \vdash \Delta} \vee_L$$

$$\frac{\Gamma \vdash A, B, \Delta}{\Gamma \vdash A \vee B, \Delta} \vee_R$$

$$\frac{\Gamma \vdash A, \Delta \quad \Gamma, B \vdash \Delta}{\Gamma, A \Rightarrow B \vdash \Delta} \Rightarrow_L$$

$$\frac{\Gamma, A \vdash B, \Delta}{\Gamma \vdash A \Rightarrow B, \Delta} \Rightarrow_R$$

$$\frac{\Gamma \vdash A, \Delta}{\Gamma, \neg A \vdash \Delta} \neg_L$$

$$\frac{\Gamma, A \vdash \Delta}{\Gamma \vdash \neg A, \Delta} \neg_R$$

$$\frac{\Gamma, A[c/x] \vdash \Delta}{\Gamma, \exists x A \vdash \Delta} \exists_L$$

$$\frac{\Gamma \vdash A[t/x], \Delta}{\Gamma \vdash \exists x A, \Delta} \exists_R$$

$$\frac{\Gamma, A[t/x] \vdash \Delta}{\Gamma, \forall x A \vdash \Delta} \forall_L$$

$$\frac{\Gamma \vdash A[c/x], \Delta}{\Gamma \vdash \forall x A, \Delta} \forall_R$$

The Intuitionistic Sequent Calculus (LJ)

$$\frac{}{\Gamma, A \vdash A} \text{ax}$$

$$\frac{\Gamma, A, B \vdash \Delta}{\Gamma, A \wedge B \vdash \Delta} \wedge_L$$

$$\frac{\Gamma \vdash A \quad \Gamma \vdash B}{\Gamma \vdash A \wedge B} \wedge_R$$

$$\frac{\Gamma, A \vdash \Delta \quad \Gamma, B \vdash \Delta}{\Gamma, A \vee B \vdash \Delta} \vee_L$$

$$\frac{\Gamma \vdash A}{\Gamma \vdash A \vee B} \vee_{R1} \quad \frac{\Gamma \vdash B}{\Gamma \vdash A \vee B} \vee_{R2}$$

$$\frac{\Gamma \vdash A \quad \Gamma, B \vdash \Delta}{\Gamma, A \Rightarrow B \vdash \Delta} \Rightarrow_L$$

$$\frac{\Gamma, A \vdash B}{\Gamma \vdash A \Rightarrow B} \Rightarrow_R$$

$$\frac{\Gamma \vdash A}{\Gamma, \neg A \vdash \Delta} \neg_L$$

$$\frac{\Gamma, A \vdash}{\Gamma \vdash \neg A} \neg_R$$

$$\frac{\Gamma, A[c/x] \vdash \Delta}{\Gamma, \exists x A \vdash \Delta} \exists_L$$

$$\frac{\Gamma \vdash A[t/x]}{\Gamma \vdash \exists x A} \exists_R$$

$$\frac{\Gamma, A[t/x] \vdash \Delta}{\Gamma, \forall x A \vdash \Delta} \forall_L$$

$$\frac{\Gamma \vdash A[c/x]}{\Gamma \vdash \forall x A} \forall_R$$

Note on Frameworks

- ▶ structural rules are not shown (contraction, weakening)
- ▶ left-rules seem **very** similar in both cases
- ▶ so, lhs formulæ can be translated by themselves
- ▶ this accounts for **polarizing** the translations
- ▶ another work [Boudard & H]:
 - ★ does not behave well in presence of cuts
 - ★ appeals to **focusing** techniques

Examples

- ▶ proofs that behave identically in classical/intuitionistic logic:

$$\frac{\overline{A, B \vdash A} \text{ ax}}{A \vdash B \Rightarrow A} \Rightarrow_R \quad \frac{\overline{A, B \vdash B} \text{ ax}}{\frac{A \wedge B \vdash B}{A \wedge B \vdash B \vee C} \wedge_L} \vee_R$$

- ▶ proof of the excluded-middle:

Classical Logic	Intuitionistic Logic
$\frac{\overline{A \vdash A} \text{ ax}}{\frac{\vdash A, \neg A}{\vdash A \vee \neg A} \neg_R} \vee_R$	

Examples

- ▶ proofs that behave identically in classical/intuitionistic logic:

$$\frac{\overline{A, B \vdash A} \text{ ax}}{A \vdash B \Rightarrow A} \Rightarrow_R \quad \frac{\overline{A, B \vdash B} \text{ ax}}{\frac{A \wedge B \vdash B}{A \wedge B \vdash B \vee C} \wedge_L} \vee_R$$

- ▶ proof of the excluded-middle:

Classical Logic	Intuitionistic Logic
$\frac{\overline{A \vdash A} \text{ ax}}{\frac{\vdash A, \neg A}{\vdash A \vee \neg A} \neg_R} \vee_R$	$\frac{??}{\vdash A \vee \neg A}$

The Excluded-Middle in Intuitionistic Logic

- is not provable. However, its **negation** is inconsistent.
- this suggests a scheme for a translation between int. and clas. logic:

$$\begin{array}{c}
 \frac{}{A \vdash A} \text{ax} \\
 \\
 \frac{}{\vdash A, \neg A} \neg R \\
 \\
 \frac{}{\vdash A \vee \neg A} \vee R
 \end{array}
 \qquad
 \begin{array}{c}
 \frac{}{A \vdash A} \text{ax} \\
 \frac{A \vdash A}{A \vdash A \vee \neg A} \vee R1 \\
 \frac{A \vdash A \vee \neg A}{\neg(A \vee \neg A), A \vdash} \neg L \\
 \frac{\neg(A \vee \neg A), A \vdash}{\neg(A \vee \neg A) \vdash \neg A} \neg R \\
 \frac{\neg(A \vee \neg A) \vdash \neg A}{\neg(A \vee \neg A) \vdash A \vee \neg A} \vee R2 \\
 \frac{\neg(A \vee \neg A), \neg(A \vee \neg A) \vdash}{\neg(A \vee \neg A) \vdash} \neg L \\
 \text{contr}
 \end{array}$$

- given a classical proof $\Gamma \vdash \Delta$, **store** Δ on the lhs, and translate:

$$\begin{array}{c}
 \text{rule } r \quad \frac{\Gamma \vdash A_1, \Delta \quad \Gamma \vdash A_2, \Delta}{\Gamma \vdash A, \Delta} \quad \text{Clas.} \\
 \\
 \frac{\Gamma, \neg A_1, \neg \Delta \vdash \quad \Gamma, \neg A_2, \neg \Delta \vdash}{\Gamma, \neg \Delta \vdash \neg \neg A_1 \quad \Gamma, \neg \Delta \vdash \neg \neg A_2} \neg R \\
 \frac{\Gamma, \neg \Delta \vdash A}{\Gamma, \neg \Delta, \neg A \vdash} \neg L \\
 \text{rule } r
 \end{array}$$

- need: $\neg\neg$ **everywhere** in Δ (and Γ)
- the proof of the “negation of the excluded middle” requires **duplication** (contraction), which partly explain why we allow several formulæ on the rhs in LK.

Kolmogorov's Translation

Kolmogorov's $\neg\neg$ -translation introduces $\neg\neg$ everywhere:

$$\begin{aligned} B^{Ko} &= \neg\neg B && \text{(atoms)} \\ (B \wedge C)^{Ko} &= \neg\neg(B^{Ko} \wedge C^{Ko}) \\ (B \vee C)^{Ko} &= \neg\neg(B^{Ko} \vee C^{Ko}) \\ (B \Rightarrow C)^{Ko} &= \neg\neg(B^{Ko} \Rightarrow C^{Ko}) \\ (\forall xA)^{Ko} &= \neg\neg(\forall xA^{Ko}) \\ (\exists xA)^{Ko} &= \neg\neg(\exists xA^{Ko}) \end{aligned}$$

Theorem

$\Gamma \vdash \Delta$ is provable in LK iff $\Gamma^{Ko}, \lrcorner\Delta^{Ko} \vdash$ is provable in LJ.

Antinegation

\lrcorner is an **operator**, such that:

- ▶ $\lrcorner\neg A = A$;
- ▶ $\lrcorner B = \neg B$ otherwise.

Light Kolmogorov's Translation

Moving negation from connectives to formulæ [Dowek& Werner]:

$$\begin{aligned} B^K &= B && \text{(atoms)} \\ (B \wedge C)^K &= (\neg\neg B^K \wedge \neg\neg C^K) \\ (B \vee C)^K &= (\neg\neg B^K \vee \neg\neg C^K) \\ (B \Rightarrow C)^K &= (\neg\neg B^K \Rightarrow \neg\neg C^K) \\ (\forall x A)^K &= \forall x \neg\neg A^K \\ (\exists x A)^K &= \exists x \neg\neg A^K \end{aligned}$$

Theorem

$\Gamma \vdash \Delta$ is provable in LK iff $\Gamma^K, \neg\neg\Delta^K \vdash$ is provable in LJ.

Correspondence

$$A^{Ko} = \neg\neg A^K$$

How does the Translation Work ?

Theorem

$\Gamma \vdash \Delta$ is provable in LK iff $\Gamma^K, \neg\Delta^K \vdash$ is provable in LJ.

Proof: Induction on the LK proof. \neg bounces. Example: rule \wedge_R .

$$\wedge_R \frac{\frac{\pi_1}{\Gamma \vdash A, \Delta} \quad \frac{\pi_2}{\Gamma \vdash B, \Delta}}{\Gamma \vdash A \wedge B, \Delta}$$

is turned into:

$$\frac{\frac{\pi'_1}{\Gamma^K, \neg A^K, \neg\Delta^K \vdash} \quad \frac{\pi'_2}{\Gamma^K, \neg B^K, \neg\Delta^K \vdash}}{\Gamma^K, \neg(\neg\neg A^K \wedge \neg\neg B^K), \neg\Delta^K \vdash} \wedge_R$$

How does the Translation Work ?

Theorem

$\Gamma \vdash \Delta$ is provable in LK iff $\Gamma^K, \neg\Delta^K \vdash$ is provable in LJ.

Proof: Induction on the LK proof. \neg bounces. Example: rule \wedge_R .

$$\wedge_R \frac{\frac{\pi_1}{\Gamma \vdash A, \Delta} \quad \frac{\pi_2}{\Gamma \vdash B, \Delta}}{\Gamma \vdash A \wedge B, \Delta}$$

is turned into:

$$\frac{\frac{\pi'_1}{\Gamma^K, \neg A^K, \neg\Delta^K \vdash} \quad \frac{\pi'_2}{\Gamma^K, \neg B^K, \neg\Delta^K \vdash}}{\Gamma^K, \neg(\neg\neg A^K \wedge \neg\neg B^K), \neg\Delta^K \vdash} \wedge_R$$

How does the Translation Work ?

Theorem

$\Gamma \vdash \Delta$ is provable in LK iff $\Gamma^K, \neg\Delta^K \vdash$ is provable in LJ.

Proof: Induction on the LK proof. \neg bounces. Example: rule \wedge_R .

$$\wedge_R \frac{\frac{\pi_1}{\Gamma \vdash A, \Delta} \quad \frac{\pi_2}{\Gamma \vdash B, \Delta}}{\Gamma \vdash A \wedge B, \Delta}$$

is turned into:

$$\neg_L \frac{\frac{\frac{\pi'_1}{\Gamma^K, \neg A^K, \neg\Delta^K \vdash} \quad \frac{\pi'_2}{\Gamma^K, \neg B^K, \neg\Delta^K \vdash}}{\Gamma^K, \neg\Delta^K \vdash \neg\neg A^K \wedge \neg\neg B^K} \quad \wedge_R}{\Gamma^K, \neg(\neg\neg A^K \wedge \neg\neg B^K), \neg\Delta^K \vdash}$$

How does the Translation Work ?

Theorem

$\Gamma \vdash \Delta$ is provable in LK iff $\Gamma^K, \neg\Delta^K \vdash$ is provable in LJ.

Proof: Induction on the LK proof. \neg bounces. Example: rule \wedge_R .

$$\wedge_R \frac{\frac{\pi_1}{\Gamma \vdash A, \Delta} \quad \frac{\pi_2}{\Gamma \vdash B, \Delta}}{\Gamma \vdash A \wedge B, \Delta}$$

is turned into:

$$\neg_R \frac{\frac{\frac{\pi'_1}{\Gamma^K, \neg A^K, \neg\Delta^K \vdash} \quad \frac{\pi'_2}{\Gamma^K, \neg B^K, \neg\Delta^K \vdash}}{\Gamma^K, \neg\Delta^K \vdash \neg\neg A^K} \quad \frac{\Gamma^K, \neg\Delta^K \vdash \neg\neg B^K}{\Gamma^K, \neg\Delta^K \vdash \neg\neg A^K \wedge \neg\neg B^K} \neg_R}{\Gamma^K, \neg(\neg\neg A^K \wedge \neg\neg B^K), \neg\Delta^K \vdash} \neg_L \wedge_R$$

Are they morphisms ?

Consider Kolmogorov's translation:

- ▶ let:

$$\begin{aligned}B \wedge_c C &= \neg\neg(B \wedge_i C) \\B \vee_c C &= \neg\neg(B \vee_i C) \\B \Rightarrow_c C &= \neg\neg(B \Rightarrow_i C) \\ \forall_c xA &= \neg\neg(\forall_i xA) \\ \exists_c xA &= \neg\neg(\exists_i xA)\end{aligned}$$

- ▶ unfortunately:

$$\begin{aligned}B^{Ko} &= \neg\neg B && \text{(atoms)} \\(B \wedge C)^{Ko} &= B^{Ko} \wedge_c C^{Ko} \\(B \vee C)^{Ko} &= B^{Ko} \vee_c C^{Ko} \\(B \Rightarrow C)^{Ko} &= B^{Ko} \Rightarrow_c C^{Ko} \\(\forall xA)^{Ko} &= \forall_c xA^{Ko} \\(\exists xA)^{Ko} &= \exists_c xA^{Ko}\end{aligned}$$

- ▶ this is not a morphism.

Are they morphisms ?

- ▶ No !

- ★ in the case of K_0 :

$$B^{K_0} = \neg\neg B(\text{atoms})$$

- ★ in the case of K :

Theorem

$\Gamma \vdash \Delta$ is provable in LK iff $\Gamma^K, \neg\Delta^K \vdash$ is provable in LJ.

- ★ exercise: these negations are necessary (hint: consider the excluded-middle and its derivatives)
- ▶ can we be more clever ?
 - ★ some **intuitionistic** right-rules are the same as **classical** right-rules. For instance, \wedge_R :

$$\frac{\Gamma \vdash A, \Delta \quad \Gamma \vdash B, \Delta}{\Gamma \vdash A \wedge B, \Delta}$$

- ★ Translate them by themselves. Gödel-Getzen translation.

Gödel-Gentzen Translation

In this translation, disjunctions and existential quantifiers are replaced by a combination of negation and their De Morgan duals:

$$\begin{aligned}B^{gg} &= \neg\neg B \\(A \wedge B)^{gg} &= A^{gg} \wedge B^{gg} \\(A \vee B)^{gg} &= \neg(\neg A^{gg} \wedge \neg B^{gg}) \\(A \Rightarrow B)^{gg} &= A^{gg} \Rightarrow B^{gg} \\(\forall xA)^{gg} &= \forall xA^{gg} \\(\exists xA)^{gg} &= \neg\forall x\neg A^{gg}\end{aligned}$$

Example of translation

$((A \vee B) \Rightarrow C)^{gg}$ is $(\neg(\neg\neg\neg A \wedge \neg\neg\neg B)) \Rightarrow \neg\neg C$

Theorem

$\Gamma \vdash \Delta$ is provable in LK iff $\Gamma^{gg}, \lrcorner\Delta^{gg} \vdash$ is provable in LJ.

Are they morphisms ?

- ▶ No !

- ★ in the case of K_0 :

$$B^{K_0} = \neg\neg B(\text{atoms})$$

- ★ in the case of K :

Theorem

$\Gamma \vdash \Delta$ is provable in LK iff $\Gamma^K, \neg\Delta^K \vdash$ is provable in LJ.

- ★ exercise: show that those negations are necessary (hint: consider the excluded-middle and its derivatives)
- ▶ can we be more clever ?
 - ★ some **intuitionistic** right-rules are the same as **classical** right-rules. For instance, \wedge_R :

$$\frac{\Gamma \vdash A, \Delta \quad \Gamma \vdash B, \Delta}{\Gamma \vdash A \wedge B, \Delta}$$

- ★ Gödel-Getzen translation:
 - ★ **is still not a morphism !**
- ▶ etc. for all the other known translations (Krivine, Kuroda)

How to make a morphism: an analysis

- ▶ Translation of, say, $A \wedge B$:

Kolmogorov	Light Kolmogorov
$\neg\neg(A^{Ko} \wedge B^{Ko})$	$(\neg\neg A^{Ko}) \wedge (\neg\neg B^{Ko})$

- ▶ Feature, double-negation:

Kolmogorov	Light Kolmogorov
on top of the connective	inside the connective

- ▶ Analysis, problem appearing in:

	Kolmogorov	Light Kolmogorov
Problem	atoms: $\neg\neg P$	statement: $\Gamma^K, \neg\Delta^K \vdash$
Solution	statement: $\Gamma^{Ko}, \neg\Delta^{Ko} \vdash$	atoms: P

- ▶ Solution: combine them !

Dowek's translation

$$\begin{aligned} B^D &= B & &= B & & \text{(atoms)} \\ (B \wedge C)^D &= B^D \wedge_c C^D & &= \neg\neg(\neg\neg B^D \wedge \neg\neg C^D) \\ (B \vee C)^D &= B^D \vee_c C^D & &= \neg\neg(\neg\neg B^D \vee \neg\neg C^D) \\ (B \Rightarrow C)^D &= B^D \Rightarrow_c C^D & &= \neg\neg(\neg\neg B^D \Rightarrow \neg\neg C^D) \\ (\forall x A)^D &= \forall_c x A^D & &= \neg\neg \forall x \neg\neg A^D \\ (\exists x A)^D &= \exists_c x A^D & &= \neg\neg \exists x \neg\neg A^D \end{aligned}$$

Theorem

$\Gamma \vdash \Delta$ is provable in LK iff $\Gamma^D, \lrcorner \Delta^D \vdash$ is provable in LJ.

Corollary

Assume A is not atomic. $\Gamma \vdash A$ is provable in LK iff $\Gamma^D \vdash A^D$ is provable in LJ.

Proof:

- ▶ $\lrcorner \lrcorner A^D = A^D$ (except in the atomic case) \square

The Price to Pay

- ▶ heavy: for each connective, 6 negations. $((A \vee B) \Rightarrow C)^D$ is $\neg\neg(\neg\neg\neg\neg(\neg\neg A \vee \neg\neg B) \Rightarrow \neg\neg C)$
- ▶ most of the time **useless**, except at the top and at the bottom of the formula
- ▶ remember Gödel-Gentzen's idea. Use De Morgan duals:

$$(A \vee B)^{gg} = \neg(\neg A^{gg} \vee \neg B^{gg})$$

- ▶ let us do the same, and divide by two the number of double negations.

A Light Morphism

Remember De Morgan,

$$A \vee B = \neg(\neg A \wedge \neg B)$$

$$A \wedge B = \neg(\neg A \vee \neg B)$$

$$A \Rightarrow B = \neg A \vee B$$

$$\neg\neg A = A$$

$$\neg\forall x A = \exists x \neg A$$

$$\neg\exists x A = \forall x \neg A$$

A Light Morphism

Remember De Morgan, and let

$$A \vee_c B = \neg(\neg A \wedge \neg B)$$

$$A \wedge_c B = \neg(\neg A \vee \neg B)$$

$$A \Rightarrow_c B = \neg(\neg\neg A \vee \neg B)$$

$$\neg_c A = \neg\neg\neg A$$

$$\forall_c xA = \neg\exists x\neg A$$

$$\exists_c xA = \neg\forall x\neg A$$

- ▶ this gives rise to a morphism, $(.)^\odot$ together with:

$$\top_c = \neg\neg\top$$

$$\perp_c = \neg\neg\perp$$

- ▶ and we can prove the theorem:

Theorem

$\Gamma \vdash \Delta$ is provable in LK iff $\Gamma^\odot, \lrcorner\Delta^\odot \vdash$ is provable in LJ.

Some Cases

Proof by induction on the proof of $\Gamma \vdash \Delta$.

- ▶ last rule \vee_R on some $A \vee B \in \Delta$. Remember:

$$\lrcorner(A \vee B)^\circ = \neg A^\circ \wedge \neg B^\circ$$

$$\frac{\Gamma \vdash A, B, \Delta}{\Gamma \vdash A \vee B, \Delta}$$

- ★ A and B are atomic: $\lrcorner A^\circ = \neg A$ and $\lrcorner B^\circ = \neg B$.

$$\frac{\Gamma^\circ, \neg A, \neg B, \lrcorner \Delta^\circ \vdash}{\Gamma^\circ, \neg A \wedge \neg B, \lrcorner \Delta^\circ \vdash}$$

- ★ if neither A and B are atomic, then A° and B° have a trailing \neg , and we remove it (bouncing):

$$\frac{\frac{\Gamma^\circ, \lrcorner A^\circ, \lrcorner B^\circ, \lrcorner \Delta^\circ \vdash}{\Gamma^\circ, \neg A^\circ, \neg B^\circ, \lrcorner \Delta^\circ \vdash}}{\Gamma^\circ, \neg A^\circ \wedge \neg B^\circ, \lrcorner \Delta^\circ \vdash} (\neg_R, \neg_L) \times 2$$

- ★ mixed case: mixed strategy.

Conclusion, Further Work

- ▶ logic with two kinds of connectives: \forall_i and \forall_c

$$\forall_{R1} \frac{\Gamma \vdash A}{\Gamma \vdash A \forall_i B} \quad \forall_{R2} \frac{\Gamma \vdash B}{\Gamma \vdash A \forall_i B}$$

- ▶ and we have:

If Γ, Δ, A contain only classical connectives, A non atomic, then $\Gamma \vdash A$ in LK iff $\Gamma \vdash A$. As well, $\Gamma \vdash \Delta$ in LK iff $\Gamma, \lrcorner\Delta \vdash$.

- ▶ next, lighter morphisms:
 - ★ from $\neg_c A = \neg\neg\neg A$ to $\neg_c A = \neg A$?
 - ★ from $A \Rightarrow_c B = \neg(\neg\neg A \vee \neg B)$ to $A \Rightarrow_c B = \neg(A \vee \neg B)$?
 - ★ we cannot always maintain the invariant $\Gamma, \lrcorner\Delta \vdash$.
 - ★ **Focusing** in LK to the rescue.